L Number	Hits	Search Text	DB	Time stamp
1	191	urotensin	USPAT;	2003/03/10 08:49
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
2	131449	mammal\$	USPAT;	2003/03/10 08:49
			US-PGPUB;	
			EPO; JPO;	
į			DERWENT	
3	114	urotensin and mammal\$	USPAT;	2003/03/10 08:49
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
4	39860	hypertens\$	USPAT;	2003/03/10 08:49
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
5	70	(urotensin and mammal\$) and hypertens\$	USPAT;	2003/03/10 08:54
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
6	95	urotensin adj II	USPAT;	2003/03/10 08:54
		•	US-PGPUB;	
			EPO; JPO;	
			DERWENT	
7	61	hypertens\$ and (urotensin adj II)	USPAT;	2003/03/10 09:05
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	
8	11949	neurodegen\$	USPAT;	2003/03/10 09:05
			US-PGPUB;	
ļ			EPO; JPO;	
			DERWENT	
9	39	(urotensin adj II) and neurodegen\$	USPAT;	2003/03/10 09:05
			US-PGPUB;	
			EPO; JPO;	
			DERWENT	

(FILE 'HOME' ENTERED AT 09:12:47 ON 10 MAR 2003)

FILE 'MEDLINE, BIOSIS, CAPLUS, EMBASE, CANCERLIT' ENTERED AT 09:13:04 ON 10 MAR 2003

```
879 S UROTENSIN II
L1
L2
            6 S NEURODGEN?
         52983 S NEURODEGEN?
L3
L4
            1 S L1 AND L3
       2947506 S NEURO?
L5
L6
           379 S L1 AND L5
      7240298 S DISEASE
L7
L8
       236783 S TRAUMA
L9
            37 S L6 AND L7
L10
            1 S L6 AND L8
            31 DUP REM L9 (6 DUPLICATES REMOVED)
L11
     1061050 S ANTAGONIST?
L12
L13
           113 S L1 AND L12
L14
           67 DUP REM L13 (46 DUPLICATES REMOVED)
L15
            5 S L14 NOT PY>1998
```

.

🚵 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
Hosted by No	CSC US Mirror s	ites: Canada China Korea T	aiwan USA	
Search Swiss-Prot/TrE	MBL	▼ for	Go Clear	

 $\mathbb{Q}9\mathbb{Q}\mathbb{Z}\mathbb{Q}3$

Printer-friendly view

Quick BlastP search

[General] [Name and origin] [References] [Comments] [Cross-references] [Keywords] [Features] [Sequence] [Tools]

Note: most headings are clickable, even if they don't appear as links. They link to the user manual or other documents.

General information about the entry					
Entry name	UR2_MOUSE				
Primary accession number	Q9QZQ3				
Secondary accession numbers	None				
Entered in Swiss-Prot in	Release 40, October 2001				
Sequence was last modified in	Release 40, October 2001				
Annotations were last modified in	Release 41, February 2003				
Name and origin of the protein					
Protein name	Urotensin II [Precursor]				
Synonyms	U-II				
	·UII				
Gene name	UTS2				
From	Mus musculus (Mouse) [TaxID: 10090]				
Taxonomy	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia;				
<u> </u>	Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Mus.				
References					

[1] SEQUENCE FROM NUCLEIC ACID.

TISSUE=Spinal cord;

MEDLINE=99416011; PubMed=10486557; [NCBI, ExPASy, EBI, Israel, Japan]

Coulouarn Y., Jegou S., Tostivint H., Vaudry H., Lihrmann I.;

"Cloning, sequence analysis and tissue distribution of the mouse and rat urotensin II precursors."; FEBS Lett. 457:28-32(1999).

Comments

- FUNCTION: HIGHLY POTENT VASOCONSTRICTOR (BY SIMILARITY).
- SUBCELLULAR LOCATION: Secreted.
- TISSUE SPECIFICITY: BRAIN-SPECIFIC. PREDOMINANTLY EXPRESSED IN MOTONEURONS OF THE BRAINSTEM AND SPINAL CORD.
- SIMILARITY: BELONGS TO THE UROTENSIN 2 FAMILY.

Copyoleht

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See http://www.isb-sib.ch/announce/ or send an email to license@isb-sib.ch).

EMBL		AF172175; AAD55767.1; [EMBL / GenBank / DDBJ] [CoDingSequence]							
MGD		MGI:1346329; Uts2.							
GeneLynx		UTS2; Mus musculus.							
SOURCE		UTS2;	Mus muscu	lus.					
Ensembl		Q9QZQ)3; Mus mu	sculus. [Entry / Contig	<u>view]</u>				
InterPro		IPR001	483; Urotei	nsin_II.					
interrio				domain structure.					
Pfam	~	PF0208	3; Urotensi	n_II; 1.			- SARA ABRINE - AMANTANA - AMANTANA - AMANTANA - AMANTANA ABRINA		
PROSITE		PS0098	<u>34;</u> UROTE	NSIN_II; 1.	MANA	**************************************	terrers — 1 decreases where 2 and dispensive transmitted where that 2 miles he will be selected to the contract of the contrac		
ProDom		[Domai	n structure	List of seq. sharing at	least 1 domain].	•			
BLOCKS		Q9QZC	<u>)3</u> .			*			
ProtoNet		Q9QZQ	<u>)3</u> .						
ProtoMap		Q9QZQ	<u>)3</u> .						
PRESAGE		Q9QZQ3.							
DIP		Q9QZQ	<u>)3</u> .						
ModBase		Q9QZQ	<u>23</u> .						
SWISS-2DPA	GE	Get reg	ion on 2D F	'AGE.			Analysis and the second		
Regywords									
Hormone; Cle	avage o	n pair o	f basic resi	dues; Signal.			The second secon		
Peatures									
1			and the Mark of State of the State of S	ugan dagan sing aut i - diri seri seringun hini daga seriab seringun danga samawan					
Key	From	To	Length	Description			Esatura alianar		
SIGNAL	1	20	20	POTENTIAL.			Feature aligner		
PROPEP	21	1 104 84 POTENTIAL.			_				
PEPTIDE	110								
DISULFID	117	Professional Continue to blanching							

Sequence infor	mation					
	[This is the leng				C64: F96486195 ecksum on the seq	137F7F4 [This is a uence]
10 MDRVPFCCLL	20 FIGLLNPLLS	30 LPVTDTGERT	40 LQLPVLEEDA	50 LRALEELERM	60 ALLQTLRQTM	**************************************
70 GTEAGESPGE	80 AGPSTETPTP	90 RGSMRKAFAG	100 QNSNTVLSRL	110 LARTRKQHKQ	120 HGAAPECFWK	· · · · · · · · · · · · · · · · · · ·
YCI						Q9QZQ3 in <u>FASTA</u> format

<u>View entry in original Swiss-Prot format</u> <u>View entry in raw text format (no links)</u> <u>Report form for errors/updates in this Swiss-Prot entry</u>

BLAST

BLAST submission on ExPASy/SIB or at NCBI (USA)



Sequence analysis tools: <u>ProtParam, ProtScale, Compute pl/Mw, PeptideMass, PeptideCutter, Dotlet</u> (Java)



ScanProsite, MotifScan



Search the **SWISS-MODEL** Repository

ExPASy Home page Site Man Search ExPASy Contact us Swisse Profession | Hosted by NCSC US Mirror sites: Canada China Korea Taiwan USA

ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot				
Hosted by NCSC US Mirror sites: Canada China Korea Taiwan USA								
Search Swiss-Prot/TrE	MBL	▼ for	Go Clear					

Q9QZQ4

Printer-friendly view Quick BlastP search

[General] [Name and origin] [References] [Comments] [Cross-references] [Keywords] [Features] [Sequence] [Tools]

Note: most headings are clickable, even if they don't appear as links. They link to the user manual or other documents.

Entry name	UR2 RAT
Primary accession number	Q9QZQ4
Secondary accession numbers	None
Entered in Swiss-Prot in	Release 40, October 2001
Sequence was last modified in	Release 40, October 2001
Annotations were last modified in	Release 40, October 2001
Name and origin of the protein	
Protein name	Urotensin II [Precursor]
Synonyms	U-II
	UII
Gene name	·UTS2
From	Rattus norvegicus (Rat) [TaxID: 10116]
Taxonomy	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia;
• •	Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Rattus.
References	

[1] SEQUENCE FROM NUCLEIC ACID.

TISSUE=Spinal cord;

MEDLINE=99416011; PubMed=10486557; [NCBI, ExPASy, EBI, Israel, Japan]

Coulouarn Y., Jegou S., Tostivint H., Vaudry H., Lihrmann I.;

"Cloning, sequence analysis and tissue distribution of the mouse and rat urotensin II precursors."; FEBS Lett. 457:28-32(1999).

Comments

- FUNCTION: HIGHLY POTENT VASOCONSTRICTOR (BY SIMILARITY).
- SUBCELLULAR LOCATION: Secreted.
- TISSUE SPECIFICITY: Brain-specific.
- SIMILARITY: BELONGS TO THE UROTENSIN 2 FAMILY.

Copyright

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See http://www.isb-sib.ch/announce/ or send an email to license@isb-sib.ch).

EMBL	AF172174; AAD55766.1; [EMBL / GenBank / DDBJ] [CoDingSequence								
EMDL		IPR001483; Urotensin_II.							
InterPro	Graphical view of domain structure.								
Pfam	PF02083; Urotensin II; 1.								
PROSITE	PS00984; UROTENSIN II; 1.								
ProDom	[Domain structure / List of seq. sharing at least 1 domain].	The state of the s							
BLOCKS	Q9QZQ4.								
ProtoNet	<u>Q9QZQ4</u> .								
ProtoMap	<u>Q9QZQ4</u> .	от на наду. 15 гот на г. неоринарија, 13 прир (Вис Вой) ВИВВисой поваја ди д ј							
PRESAGE	Q9QZQ4.	THE RESIDENCE OF THE PROPERTY							
DIP	<u>Q9QZQ4</u> .	e. ma · mace as graves meligia are <u>also graves or presenting constitute or graves and constitute</u>							
ModBase	Q9QZQ4.								
SWISS-2DPAC	GE Get region on 2D PAGE.	COLD TO THE COMPANY OF THE COLD TO THE COL							
Keywords	The state of the s								
	avage on pair of basic residues; Signal.	The second of th							
Features		THE CHARGE DO IN THE RESIDENCE OF THE PARTY							
Benefic									
Vois	From To Length Description								
K ey SIGNAL	1 20 20 POTENTIAL.	eature aligner							
PROPEP	21 104 84 POTENTIAL.								
PEPTIDE	110 123 14 UROTENSIN II.								
DISULFID	117 122 BY SIMILARITY.	eature table viewer							
ï		•							
Sequence info	amothon.								
Length: 123 AA of the unproces	A [This is the length Molecular weight: 13614 Da [This is CRC64: E4F1A8EE12								
	నారి కామానికి చేసుకు కానీకి కల్పు కారాగులు నుండి మహారాముల హీకు <u>మముందా మేకు ముఖుముముంది. ముఖు మమీములు</u>	icej							
10	20 30 40 50 60								
MDRVPFCCLL	FVGLLNPLLS FPVTDTGEMS LQLPVLEENA LRALEELERT ALLQTLRQTV								
70	80 90 100 110 120								
 GTEAEGSLGO	ADPSAETPTP RGSLRKALTG ODSNTVLSRL LARTRKORKO HGTAPECFWK								
YCI									
ICI		Q9QZQ4 in FASTA							
		£ 424 1110111							

<u>View entry in original Swiss-Prot format</u> <u>View entry in raw text format (no links)</u>

Report form for errors/updates in this Swiss-Prot entry

BLAST

BLAST submission on ExPASy/SIB or at NCBI (USA)



Sequence analysis tools: <u>ProtParam</u>, <u>ProtScale</u>, <u>Compute</u> <u>pI/Mw</u>, <u>PeptideMass</u>, <u>PeptideCutter</u>, <u>Dotlet</u> (Java)

<u>format</u>



ScanProsite, MotifScan



Search the **SWISS-MODEL** Repository

É ExPASy Home page Site Map Search ExPASy Contact us Swiss-Prot

№ ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
Hosted by NO	CSC US Mirror s	sites: <u>Canada China Korea</u> 7	aiwan USA	
Search Swiss-Prot/TrE	MBL	▽ for	Go; Clear	

Printer-friendly view Quick BlastP search

[General] [Name and origin] [References] [Comments] [Cross-references] [Keywords] [Features] [Sequence] [Tools]

Note: most headings are clickable, even if they don't appear as links. They link to the user manual or other documents.

General information about the c	
Entry name	UR2_HUMAN
Primary accession number	O95399
Secondary accession number	Q9UKP7
Entered in Swiss-Prot in	Release 40, October 2001
Sequence was last modified in	Release 40, October 2001
Annotations were last modified in	Release 41, February 2003
Name and origin of the protein	
Protein name	Urotensin II [Precursor]
Synonyms	U-II
	UII
Gene name	UTS2
From	Homo sapiens (Human) [TaxID: 9606]
Taxonomy	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia;
	Eutheria; Primates; Catarrhini; Hominidae; Homo.
References	

[1] SEQUENCE FROM NUCLEIC ACID.

TISSUE=Spinal cord:

MEDLINE=99080095; PubMed=9861051; [NCBI, ExPASy, EBI, Israel, Japan]

Coulouarn Y., Lihrmann I., Jegou S., Anouar Y., Tostivint H., Beauvillain J.C., Conlon J.M., Bern H.A., Vaudry H.; "Cloning of the cDNA encoding the urotensin II precursor in frog and human reveals intense expression of the urotensin II gene in motoneurons of the spinal cord.";

Proc. Natl. Acad. Sci. U.S.A. 95:15803-15808(1998).

[2] SEQUENCE FROM NUCLEIC ACID.

MEDLINE=99427933; PubMed=10499587; [NCBI, ExPASy, EBI, Israel, Japan]

Ames R.S., Sarau H.M., Chambers J.K., Willette R.N., Aiyar N.V., Romanic A.M., Louden C.S., Foley J.J., Sauermelch C.F., Coatney R.W., Ao Z., Disa J., Holmes S.D., Stadel J.M., Martin J.D., Liu W.-S., Glover G.I., Wilson S., McNulty D.E., Ellis C.E., Elshourbagy N.A., Shabon U., Trill J.J., Hay D.W.P., Ohlstein E.H., Bergsma D.J., Douglas S.A.; "Human urotensin-II is a potent vasoconstrictor and agonist for the orphan receptor GPR14."; Nature 401:282-286(1999).

[3] SEQUENCE FROM NUCLEIC ACID.

Pearce A .:

Submitted (DEC-1999) to the EMBL/GenBank/DDBJ databases.

Comments

- FUNCTION: HIGHLY POTENT VASOCONSTRICTOR.
- SUBCELLULAR LOCATION: Secreted.
- TISSUE SPECIFICITY: Brain-specific.
- SIMILARITY: BELONGS TO THE UROTENSIN 2 FAMILY.

|Copyrelepto

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See http://www.isb-sib.ch/announce/ or send an email to https://www.isb-sib.ch/announce/ or <a href="https://www.isb-sib.ch

Cross-reference	98)	
L	AF104118; AAD13070.1;[EMBL / GenBank / DDBJ] [CoDingSequence	el
EMBL	AF140630; AAD55577.1; [EMBL / GenBank / DDBJ] [CoDingSequence	
	Z98884; CAB63148.1; [EMBL / GenBank / DDBJ] [CoDingSequence	
Genew	HGNC:12636; UTS2.	
CleanEx	HGNC:12636; UTS2.	a 1191 a american
MIM	604097 [NCBI / EBI].	
GeneCards	UTS2.	
GeneLynx	UTS2; Homo sapiens.	
SOURCE	UTS2; Homo sapiens.	
Ensembl	O95399; Homo sapiens. [Entry / Contig view]	The state of the other control of the second section of the section of t
1 / 5	IPR001483; Urotensin II.	
InterPro	Graphical view of domain structure.	
Pfam	PF02083; Urotensin_II; 1.	- THE CONTRACTOR OF THE PROPERTY OF THE PROPER
PROSITE	<u>PS00984;</u> UROTENSIN_II; 1.	
ProDom	[Domain structure / List of seq. sharing at least 1 domain].	
BLOCKS	<u>095399</u> .	
ProtoNet	<u>095399</u> .	
ProtoMap	<u>095399</u> .	
PRESAGE	<u>095399</u> .	TAR BUT PURE CONTRACTOR AND TAR A CONTRACTOR FOR THE CONTRACTOR AND
DIP	<u>095399</u> .	
ModBase	<u>095399</u> .	
SWISS-2DPAG	E Get region on 2D PAGE.	
Kaywords		
Hormone; Clea	vage on pair of basic residues; Signal.	
Features		
Key	From To Length Description	
SIGNAL	1 20 20 POTENTIAL.	
PROPEP	<u>21 110</u> 90	Feature Feature
PEPTIDE	<u>114 124</u> 11 UROTENSIN II.	<u>aligner</u>
DISULFID	118 123 BY SIMILARITY.	
CONFLICT	1 19 MYKLASCCLLFIGFLNPLL ->	P φ Feature table
	METNVFHLMLCVTSARTH KSTSLCFGHFNSYP (IN REF. 2).	<u>viewer</u>
CONFLICT	24 27 LDSR -> IHDLLL (IN REF. 2).	
		
Sequence infor		
Length: 124 AA of the unprocess	[This is the length Molecular weight: 14295 Da [This is CRC64: C7A5FC7EF]	
•		ence]
	20 30 40 50 60	
10 		
		n 1
MYKLASCCLL		· · · · · · · · · · · · · · · · · · ·
MYKLASCCLL 70	80 90 100 110 120 	. 17
MYKLASCCLL 70		n n ha a a a a a a a a a a a a a a a a a
MYKLASCCLL 70	80 90 100 110 120 	The state of the s
MYKLASCCLL 70	80 90 100 110 120 	4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
MYKLASCCLL 70	80 90 100 110 120 	· · · · · · · · · · · · · · · · · · ·
MYKLASCCLL 70 GAERGDILRK	80 90 100 110 120 	O95399 in <u>FASTA</u>
MYKLASCCLL 70 GAERGDILRK	80 90 100 110 120 	O95399 in <u>FASTA</u>

<u>View entry in original Swiss-Prot format</u> <u>View entry in raw text format (no links)</u> <u>Report form for errors/updates in this Swiss-Prot entry</u>

BLAST

BLAST submission on ExPASy/SIB or at NCBI (USA)



Sequence analysis tools: <u>ProtParam</u>, <u>ProtScale</u>, <u>Compute</u> <u>pl/Mw</u>, <u>PeptideMass</u>, <u>PeptideCutter</u>, <u>Dotlet</u> (Java)



ScanProsite, MotifScan



Search the **SWISS-MODEL** Repository

<u>ExPASy Home page</u>	Site Map	<u>Search ExPASy</u>	Contact us	<u>Swiss-Prot</u>
Hosted by NO	CSC US Mirror sites	s: Canada China Korea T	aiwan USA	

3 of 3

ExPASy Home page	Site Map	Search ExPASy	Contact us .	Swiss-Prot
Hosted by NO	CSC US Mirror s	ites: <u> Canada</u> <u>China</u> <u>Korea</u> <u>T</u>	aiwan USA	
Search Swiss-Prot/TrE	MBL	▼ for '	Go Clear]

P49684

Printer-friendly view Quick BlastP search

[General] [Name and origin] [References] [Comments] [Cross-references] [Keywords] [Features] [Sequence] [Tools]

Note: most headings are clickable, even if they don't appear as links. They link to the <u>user manual</u> or <u>other documents</u>.

General information about the c	
Entry name	UR2R_RAT
Primary accession number	P49684
Secondary accession number	P48041
Entered in Swiss-Prot in	Release 33, February 1996
Sequence was last modified in	Release 38, July 1999
Annotations were last modified in	Release 41, February 2003
Name and origin of the protein	
Protein name	Urotensin II receptor
Synonyms	UR-II-R
	G protein-coupled sensory epithelial neuropeptide-like receptor SENR
Gene name	GPR14
From	Rattus norvegicus (Rat) [TaxID: 10116]
Taxonomy	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Rattus.
References	

Recerences

[1] SEQUENCE FROM NUCLEIC ACID.

MEDLINE=96115583; PubMed=8666380; [NCBI, ExPASy, EBI, Israel, Japan]

Marchese A., Heiber M., Nguyen T., Heng H.H.Q., Saldivia V.R., Cheng R., Murphy P.M., Tsui L.-C., Shi X., Gregor P., George S.R., O'Dowd B.F., Docherty J.M.;

"Cloning and chromosomal mapping of three novel genes, GPR9, GPR10, and GPR14, encoding receptors related to interleukin 8, neuropeptide Y, and somatostatin receptors."; Genomics 29:335-344(1995).

[2] SEQUENCE FROM NUCLEIC ACID.

TISSUE=Circumvallate papillae;

STRAIN=Sprague-Dawley;

MEDLINE=95251679; PubMed=7733947; [NCBI, ExPASy, EBI, Israel, Japan]

Tal M., Ammar D.A., Karpuj M., Krizhanovsky V., Naim M., Thompson D.A.;

"A novel putative neuropeptide receptor expressed in neural tissue, including sensory epithelia."; Biochem. Biophys. Res. Commun. 209:752-759(1995).

[3] SEQUENCE FROM NUCLEIC ACID.

TISSUE=Urinary bladder;

STRAIN=Wistar;

Suga H., Takao K.;

"Expression of the rat SENR in the urinary bladder tissues.";

Submitted (MAR-1998) to the EMBL/GenBank/DDBJ databases.

[4] SEQUENCE FROM NUCLEIC ACID.

TISSUE=Pheochromocytoma;

Liu H., Zou M., Suga H., Takao K.;

"The SENR/GPR14 expresses in rat pheochromocytoma PC 12 cells.";

Submitted (JUL-1999) to the EMBL/GenBank/DDBJ databases.

Comments

- FUNCTION: HIGH AFINITY RECEPTOR FOR UROTENSIN II. THE ACTIVITY OF THIS RECEPTOR IS MEDIATED BY A G-PROTEIN THAT ACTIVATE A PHOSPHATIDYLINOSITOL-CALCIUM SECOND MESSENGER SYSTEM (BY SIMILARITY).
- SUBCELLULAR LOCATION: Integral membrane protein.
- TISSUE SPECIFICITY: PREFERENTIALLY EXPRESSED IN NEURAL AND SENSORY TISSUES.
- SIMILARITY: BELONGS TO FAMILY 1 OF G-PROTEIN COUPLED RECEPTORS.


Copyulant

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See http://www.isb-sib.ch/announce/ or send an email to license@isb-sib.ch).

ilcense(@iso-sib.ch).	
Cross-references	
EMBL	U32673; AAC52593.1; [EMBL / GenBank / DDBJ] [CoDingSequence] U23483; AAA80111.1; [EMBL / GenBank / DDBJ] [CoDingSequence] AB012210; BAA25251.1; [EMBL / GenBank / DDBJ] [CoDingSequence] AB029611; BAA82357.1; [EMBL / GenBank / DDBJ] [CoDingSequence]
InterPro	IPR000276; GPCR_Rhodpsn. Graphical view of domain structure.
Pfam	<u>PF00001;</u> 7tm_1; 1.
PRINTS	PR00237; GPCRRHODOPSN.
PROSITE	<u>PS00237;</u> G_PROTEIN_RECEP_F1_1; 1. <u>PS50262;</u> G_PROTEIN_RECEP_F1_2; 1.
GPCRDB	<u>P49684</u> ; UR2R_RAT.
GPCRDB-Snakes	<u>P49684</u> .
ProDom	[Domain structure / List of seq. sharing at least 1 domain].
BLOCKS	<u>P49684</u> .
ProtoNet	<u>P49684</u> .
ProtoMap	<u>P49684</u> .
PRESAGE	<u>P49684</u> .
DIP	<u>P49684</u> .
ModBase	<u>P49684</u> .
SWISS-2DPAGE	Get region on 2D PAGE.
Keywords	
G-protein coupled r	eceptor; Transmembrane; Glycoprotein.
Features	

Key	From	То	Length	Description		
DOMAIN	1	54	54	EXTRACELLULAR (POTENTIAL).	*	•
TRANSMEM	55	77	23	1 (POTENTIAL).		
DOMAIN	78	87	10	CYTOPLASMIC (POTENTIAL).	1	
TRANSMEM	88	113	26	2 (POTENTIAL).		
DOMAIN	114	124	11	EXTRACELLULAR (POTENTIAL).		
TRANSMEM	125	146	22	3 (POTENTIAL).		
DOMAIN	147	167	21	CYTOPLASMIC (POTENTIAL).	-	
TRANSMEM	168	186	19	4 (POTENTIAL).		
DOMAIN	187	209	23	EXTRACELLULAR (POTENTIAL).		Feature aligner
TRANSMEM	210	232	23	5 (POTENTIAL).		
DOMAIN	233	258	26	CYTOPLASMIC (POTENTIAL).		T 11
TRANSMEM	259	284	26	6 (POTENTIAL).	\$,0	Feature table
DOMAIN	285	299	15	EXTRACELLULAR (POTENTIAL).	400	<u>viewer</u>
TRANSMEM	300	321	22	7 (POTENTIAL).	į	
DOMAIN	322	386	65	CYTOPLASMIC (POTENTIAL).	1	
CARBOHYD	29	2,9		N-LINKED (GLCNAC) (POTENTIAL).		
CARBOHYD	33	33		N-LINKED (GLCNAC) (POTENTIAL).		
DISULFID	123	199		BY SIMILARITY.		
CONFLICT	315	315		F -> L (IN REF. <u>1</u>).		

Sequence information



View entry in original Swiss-Prot format
View entry in raw text format (no links)
Report form for errors/updates in this Swiss-Prot entry

BLAST

BLAST submission on ExPASy/SIB or at NCBI (USA)



Sequence analysis tools: <u>ProtParam</u>, <u>ProtScale</u>, <u>Compute</u> <u>pl/Mw</u>, <u>PeptideMass</u>, <u>PeptideCutter</u>, <u>Dotlet</u> (Java)



ScanProsite, MotifScan



Search the **SWISS-MODEL** Repository

ExPASy Home page Site Map Search ExPASy Contact us Swiss-Prot

Hosted by NCSC US Mirror sites: Canada China Korea Taiwan USA

& ExPASy Home	e page	Site Map	Search	ExPASy	Contact us	Swiss-Prot
	Hosted by NC	SC US Mirror	sites: Canada	China Korea	Taiwan USA	
Search	Swiss-Prot/TrEM	BL	▼ for		Go Clear	7

Q9UKP6

Printer-friendly view

Quick BlastP search

[General] [Name and origin] [References] [Comments] [Cross-references] [Keywords] [Features] [Sequence] [Tools]

Note: most headings are clickable, even if they don't appear as links. They link to the user manual or other documents.

General information about the c	mtry
Entry name	UR2R_HUMAN
Primary accession number	Q9UKP6
Secondary accession numbers	None
Entered in Swiss-Prot in	Release 40, October 2001
Sequence was last modified in	Release 40, October 2001
Annotations were last modified in	Release 40, October 2001
Name and origin of the protein	
Protein name	Urotensin II receptor
Synonym	UR-II-R
Gene name	GPR14
From	Homo sapiens (Human) [TaxID: 9606]
Taxonomy	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
Deferrances	1 1 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7

References

[1] SEQUENCE FROM NUCLEIC ACID.

MEDLINE=99427933; PubMed=10499587; [NCBI, ExPASy, EBI, Israel, Japan]

Ames R.S., Sarau H.M., Chambers J.K., Willette R.N., Aiyar N.V., Romanic A.M., Louden C.S., Foley J.J., Sauermelch C.F., Coatney R.W., Ao Z., Disa J., Holmes S.D., Stadel J.M., Martin J.D., Liu W.-S., Glover G.I., Wilson S., McNulty D.E., Ellis C.E., Elshourbagy N.A., Shabon U., Trill J.J., Hay D.W.P., Ohlstein E.H., Bergsma D.J., Douglas S.A.; "Human urotensin-II is a potent vasoconstrictor and agonist for the orphan receptor GPR14."; Nature 401:282-286(1999).

Comments


- FUNCTION: HIGH AFINITY RECEPTOR FOR UROTENSIN II. THE ACTIVITY OF THIS RECEPTOR IS MEDIATED BY A G-PROTEIN THAT ACTIVATE A PHOSPHATIDYLINOSITOL-CALCIUM SECOND MESSENGER SYSTEM.
- SUBCELLULAR LOCATION: Integral membrane protein.
- TISSUE SPECIFICITY: MOST ABUNDANT EXPRESSION IN THE HEART AND PANCREAS.
- SIMILARITY: BELONGS TO FAMILY 1 OF G-PROTEIN COUPLED RECEPTORS.

Conxelebt

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See http://www.isb-sib.ch/announce/ or send an email to license@isb-sib.ch).

EMBL		AF140	631; AAD5	55578.1; [EMBL / GenBank / DDBJ] [CoDingSequence]
Genew		HGNC	:4468; GPR	114.
CleanEx			:4468; GPR	
MIM			NCBI / EI	the control of the co
GeneCards		GPR14		<u> </u>
GeneLynx			; Homo sap	ione
SOURCE				and the control of th
			; Homo sap	
Ensembl				apiens. [Entry / Contig view]
InterPro				R_Rhodpsn.
Pfam			01; 7tm 1; 1	domain structure.
PRINTS			;	HODOPSN.
PROSITE	11. 1. 1. 10m 11	PS0023	7; G_PRO	TEIN_RECEP_F1_1; 1. TEIN_RECEP_F1_2; 1.
GPCRDB				The state of the s
			<u> 26;</u> UR2R_E	IIUMAN,
GPCRDB-Snal	kes	Q9UKI		
ProDom		_		/ List of seq. sharing at least 1 domain].
BLOCKS		Q9UKI		The state of the s
ProtoNet		Q9UKI		n to the time to the transition of the second of the transition and the transition of the transition o
ProtoMap		Q9UKI		
PRESAGE		Q9UKI	<u>°6</u> .	
DIP		Q9UKI	<u>°6</u> .	
ModBase		Q9UKI	<u>26</u> .	
SWISS-2DPAG	GE	Get reg	ion on 2D P	PAGE.
Keywords				
G-protein cou	pled rec	eptor;	<u>Fransmemt</u>	brane; Glycoprotein.
Features				
Key	From	To	Length	Description
DOMAIN	1	54	54	EXTRACELLULAR (POTENTIAL).
TRANSMEM	55	77	23	1 (POTENTIAL).
DOMAIN	78	87	10	CYTOPLASMIC (POTENTIAL).
TRANSMEM	88	113	26	2 (POTENTIAL).
DOMAIN	114	124	11	EXTRACELLULAR (POTENTIAL).
TRANSMEM	125	146	22	3 (POTENTIAL).
DOMAIN	147	167	21	CYTOPLASMIC (POTENTIAL).
TRANSMEM	168	186	19	4 (POTENTIAL). Feature aligner
DOMAIN	187	209	23	EXTRACELLULAR (POTENTIAL).
TRANSMEM	210	232	23	5 (POTENTIAL).
DOMAIN	233	258	26	CYTOPLASMIC (POTENTIAL). © Feature table
TRANSMEM	259	284	26	6 (POTENTIAL). viewer
DOMAIN	285	297	13	EXTRACELLULAR (POTENTIAL).
TRANSMEM	298	318	21	7 (POTENTIAL).
DOMAIN	319	389	71	CYTOPLASMIC (POTENTIAL).
CARBOHYD	29	29		N-LINKED (GLCNAC) (POTENTIAL).
CARBOHYD	33	33		N-LINKED (GLCNAC) (POTENTIAL).
DISULFID	123	199		BY SIMILARITY.
Sequence info	rodions			

Length: 389 AA Molecular weight: 42130 Da CRC64: 6D6A88DBF78400CE [This is a checksum on the sequence]



<u>View entry in original Swiss-Prot format</u> <u>View entry in raw text format (no links)</u>

Report form for errors/updates in this Swiss-Prot entry

BLAST

BLAST submission on ExPASy/SIB or at NCBI (USA)



Sequence analysis tools: <u>ProtParam</u>, <u>ProtScale</u>, <u>Compute pl/Mw</u>, <u>PeptideMass</u>, <u>PeptideCutter</u>, <u>Dotlet</u> (Java)



ScanProsite, MotifScan



Search the **SWISS-MODEL** Repository

the state of the s		and the second s	Committee of the Commit	Filter Michigan agripm, when in Proceedings are a supersymptotic processing to a
📤 ExPASy Home page	<u>Site Map</u>	Search ExPASy	Contact us	<u>Swiss-Prot</u>
Hosted by NO	CSC US Mirror site	s: Canada China Korea T	aiwan USA	

ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
Hosted by No	CSC US Mirror s	ites: Canada China Korea	Taiwan USA	
Search Swiss-Prot/TrE	MBL	▼ for	Go Clear	

P49220

Printer-friendly view Quick BlastP search

[General] [Name and origin] [References] [Comments] [Cross-references] [Keywords] [Features] [Sequence] [Tools]

Note: most headings are clickable, even if they don't appear as links. They link to the <u>user manual</u> or <u>other documents</u>.

General information about the c	miry
Entry name	UR2R_BOVIN
Primary accession number	P49220
Secondary accession numbers	None
Entered in Swiss-Prot in	Release 33, February 1996
Sequence was last modified in	Release 33, February 1996
Annotations were last modified in	Release 40, October 2001
Name and origin of the protein	
Protein name	Urotensin II receptor [Fragment]
Synonyms	UR-II-R
:	G protein-coupled sensory epithelial neuropeptide-like receptor SENR
Gene name	GPR14
From	Bos taurus (Bovine) [TaxID: 9913]
Taxonomy	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea; Bovidae; Bovinae; Bos.
Pelerances	

References

[1] SEQUENCE FROM NUCLEIC ACID.

TISSUE=Retina;

MEDLINE=95251679; PubMed=7733947; [NCBI, ExPASy, EBI, Israel, Japan]

Tal M., Ammar D.A., Karpuj M., Krizhanovsky V., Naim M., Thompson D.A.;

"A novel putative neuropeptide receptor expressed in neural tissue, including sensory epithelia.";

Biochem. Biophys. Res. Commun. 209:752-759(1995).

Comments

- FUNCTION: HIGH AFINITY RECEPTOR FOR UROTENSIN II. THE ACTIVITY OF THIS RECEPTOR IS MEDIATED BY A G-PROTEIN THAT ACTIVATE A PHOSPHATIDYLINOSITOL-CALCIUM SECOND MESSENGER SYSTEM (BY SIMILARITY).
- SUBCELLULAR LOCATION: Integral membrane protein.
- TISSUE SPECIFICITY: EXPRESSED IN NEURAL TISSUE, INCLUDING SENSORY EPITHELIA.
- SIMILARITY: BELONGS TO FAMILY 1 OF G-PROTEIN COUPLED RECEPTORS.

Copyuight

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See http://www.isb-sib.ch/announce/ or send an email to license@isb-sib.ch).

EMBL	U2345	9; AAC48464.1;	[EMBL / GenI	Bank / DDBJ] [C	oDingSequence]				
		IPR000276; GPCR Rhodpsn.							
InterPro		cal view of doma							
Pfam	PF0000	<u>PF00001;</u> 7tm_1; 1.							
PRINTS		PR00237; GPCRRHODOPSN.							
DD OCUME	PS0023	PS00237; G PROTEIN RECEP F1 1; PARTIAL.							
PROSITE		PS50262; G_PROTEIN_RECEP_F1_2; 1.							
BLOCKS	P49220	<u>.</u> .							
ProtoNet	P49220	<u>)</u> .							
ProtoMap	P49220	<u>)</u> .							
PRESAGE	P49220	<u>)</u> .							
DIP	P49220	<u>).</u>	TO SERVICE OF THE PARTY OF THE	entern van een een een een een een een een een e	THE AND ESTABLISHED AND AS ESTABLISHED THE STREET ASSESSMENT AS	man i remand pi anno morano del mando dela mando del mando del mando del mando del mando del mando del man			
ModBase	P49220	<u>).</u>	· · · · · · · · · · · · · · · · · · ·						
SWISS-2DPAG	E Get reg	ion on 2D PAGE	<u>2</u> .						
Keywords	-		· -			A THE STATE OF THE			
G-protein coup	oled receptor; I	`ransmembrane	; <u>Glycoprotein</u> .						
Features									
Key	From To	Length Des	cription						
NON_TER	1 1								
TRANSMEM	<1 16	>16 2	(POTENTIAL) .						
DOMAIN	<u>17 27</u>	11 EXT	RACELLULAR	(POTENTIAL)					
TRANSMEM	28 49	22 3	(POTENTIAL).						
DOMAIN	50 70	21 CYT	OPLASMIC (F	POTENTIAL).		Feature aligner			
TRANSMEM	71 89		(POTENTIAL) .						
DOMAIN	90 111		RACELLULAR	(POTENTIAL)		- -			
TRANSMEM	<u>112 135</u>		(POTENTIAL) .		<u> </u>	Feature table viewer			
DOMAIN	<u>136 161</u>		COPLASMIC (F			reature table viewer			
TRANSMEM	<u>162 187</u>		(POTENTIAL).		<u> </u>	4			
DOMAIN	188 194		RACELLULAR		•				
TRANSMEM	195 215		(POTENTIAL).						
DISULFID	<u>26 102</u>	POT	TENTIAL.						
NON_TER	215 215	<u> </u>			<u> </u>				
Sequence infor									
Length: 215 AA length of the par			ight: 24575 Da [e partial sequenc		: 97F2D5041D9 im on the sequen	6A624 [This is a ce]			
10	20	30	40	50	60	The second secon			
' I	İ			1)			
LLYLLSIPFI	VATYVTKRWH	FGDVGCRVLF	SLDFLTMHAS	IFTLTLMSRE	RYAAVVRPLD				
70 	80 	90 	100	110	120				
TVQRSKGYRK	VLALGTWLLA	LLLALPMMLA	IRLVRRGHKS	LCLPAWGQRT	HRAYLTLLFG	The state of the s			
130	140	150	160	170	180				
 TSIVGPGVVI	 GLLYVRLARA	 YWLSQRASFT	 QTRRLPNPRV	LYLILGIVLL	FWACFLPFWL				
190	200	210							
į	1	1							
WQLLAQYRGA	PPLAPRSARI	VNYLTTCLTY	GNSCV			D40220 in EASTA			
						P49220 in <u>FASTA</u> format			

<u>View entry in original Swiss-Prot format</u> <u>View entry in raw text format (no links)</u>

Report form for errors/updates in this Swiss-Prot entry

BLAST

BLAST submission on ExPASy/SIB or at NCBI (USA)



Sequence analysis tools: <u>ProtParam</u>, <u>ProtScale</u>, <u>Compute</u> <u>pI/Mw</u>, <u>PeptideMass</u>, <u>PeptideCutter</u>, <u>Dotlet</u> (Java)



ScanProsite, MotifScan



Search the **SWISS-MODEL** Repository

ExPASy Home page Site Map Search ExPASy Contact us Swiss-Prot

Hosted by NCSC US Mirror sites: Canada China Korea Taiwan USA